

DR. MICHELLE A KESKE (Orcid ID : 0000-0003-4214-7628)

DR. DINO PREMILOVAC (Orcid ID : 0000-0003-2770-4713)

Article type : Symposium Paper - AuPS

### **Metabolic-vascular coupling in skeletal muscle: a potential role for capillary pericytes?**

Emily Attrill<sup>1</sup>, Ciaran Ramsay<sup>1</sup>, Renee Ross<sup>1</sup>, Stephen Richards<sup>1</sup>, Brad A Sutherland<sup>1</sup>, Michelle A Keske<sup>2</sup>, Etto Eringa<sup>3</sup> and Dino Premilovac<sup>1</sup>

#### **Author affiliations**

<sup>1</sup> School of Medicine, College of Health and Medicine, University of Tasmania, Hobart, Tasmania, Australia

<sup>2</sup> The Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition Sciences, Deakin University, Melbourne, Victoria, Australia

<sup>3</sup> Laboratory for Physiology, VU University Medical Center, Amsterdam, the Netherlands.

**Short title:** Pericyte control of muscle capillary perfusion

#### **Corresponding Author:**

Dino Premilovac (PhD)

School of Medicine, College of Health and Medicine, University of Tasmania

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](https://doi.org/10.1111/1440-1681.13208). Please cite this article as [doi: 10.1111/1440-1681.13208](https://doi.org/10.1111/1440-1681.13208)

This article is protected by copyright. All rights reserved

17 Liverpool Street, Hobart, TAS 7000, Australia

Phone: +613 6226 2701

Email: Dino.Premilovac@utas.edu.au

## **ABSTRACT**

The matching of capillary blood flow to metabolic rate of the cells within organs and tissues is a critical microvascular function which ensures appropriate delivery of hormones and nutrients, and the removal of waste products. This relationship is particularly important in tissues where local metabolism, and hence capillary blood flow, must be regulated to avoid a mismatch between nutrient demand and supply that would compromise normal function. The consequences of a mismatch in microvascular blood flow and metabolism are acutely apparent in the brain and heart, where a sudden cessation of blood flow, for example following an embolism, acutely manifests as stroke or myocardial infarction. Even in more resilient tissues such as skeletal muscle, a short-term mismatch reduces muscle performance and exercise tolerance, and can cause intermittent claudication. In the longer-term, a microvascular-metabolic mismatch in skeletal muscle reduces insulin-mediated muscle glucose uptake, leading to disturbances in whole body metabolic homeostasis. While the notion that capillary blood flow is fine tuned to meet cellular metabolism is well-accepted, the mechanisms that control this function and where and how different parts of the vascular tree contribute to capillary blood flow regulation remains poorly understood. Here, we discuss the emerging evidence implicating pericytes, mural cells that surround capillaries, as key mediators that match tissue metabolic demand with adequate capillary blood flow in a number of organs, including skeletal muscle.

**KEY WORDS:** Pericytes, capillary blood flow, microvasculature, skeletal muscle

## **1. INTRODUCTION**

### **1.1 Regulation of blood flow**

From a top-down point of view, whole body blood flow is equal to the total cardiac output. At the organ level, cardiac output is rationed between organs to meet their metabolic demands and specific functions. As an example, the brain receives a relatively constant proportion of total cardiac output (15-20%) to ensure appropriate supply of nutrients to working neurons (1). The kidneys receive approximately the same amount of blood flow to ensure adequate blood filtration and blood volume/pressure homeostasis (1). In general, total organ blood flow, is determined by the diameter of the arteriolar resistance vessels within a specific organ. The appropriate rationing of cardiac output between organs has long been proposed to be a balance between two opposing factors: (i) the sympathetic nervous system (SNS) mediating vasoconstriction to maintain mean arterial blood pressure; and (ii) organ-specific metabolic rate-mediated vasodilation and increased blood flow to ensure appropriate cellular/tissue function (2). While it is well accepted that the SNS innervates arteries and arterioles to regulate their tone and hence blood flow, it remains to be determined how increased tissue metabolism can stimulate site-specific increases in local capillary blood flow.

### **1.2 Regulation of local blood flow**

Within almost all organs, local tissue metabolic demand can vary at any given time. This is best exemplified by skeletal muscles where resting metabolism is low and hence capillary perfusion is appropriately limited to conserve whole body metabolic homeostasis. In contrast, during intense exercise, metabolic demand within contracting muscles can increase as much as 100-fold and this is matched by equal increases in capillary perfusion (2). Ever since the work of Nobel Prize winning physiologist August Krogh in the 1920s, the prevailing view has emphasized arterioles as regulators of capillary blood flow and therefore determinants of muscle nutrient supply (3). While arterioles undoubtedly play an integral role in this process, for appropriate coupling of metabolism and blood flow, the capillary network itself must be able to detect and respond to local tissue derived signals to

appropriately regulate both capillary and arteriolar perfusion. Despite the early documentation by Krogh, the physiological and biochemical mechanisms that dictate this local metabolic-vascular coupling remain to be identified and characterized.

### **1.3 Circulating versus paracrine factors in the regulation of local blood flow**

From a systemic view point, circulating factors such as angiotensin II modulate arterial tone to regulate and maintain mean arterial pressure. Importantly, angiotensin II, as well as other circulating factors such as insulin, can also have profound and specific effects within the skeletal muscle vasculature to regulate perfusion and hence muscle metabolism (4). In contrast, metabolites (e.g. adenosine) and ions (e.g.  $K^+$ ) derived from muscle metabolism are released into the extracellular space and are proposed to mediate a prompt and highly localised vascular response to increase capillary blood flow in a site-specific manner via a mechanism termed ascending vasodilation (5, 6). Championed by Steven Segal and colleagues (5-7), this involves the initiation and transmission of a hyperpolarizing wave in a cell-cell manner via gap junctions upstream to feed arteries to increase arteriolar blood flow and hence capillary perfusion (see Bagher and Segal 2011 for review) (6). However, much of what is known about these vascular integrating mechanisms was derived from local application of acetylcholine (ACh), which may activate different signaling pathways than the signals derived from muscle metabolism. Following up on these studies, others have utilised single muscle fibre stimulation to show that these conducted vasodilatory signals can originate in capillaries and propagate via gap junctions to upstream arterioles to modulate blood flow (8-10). While some evidence suggests that purinergic signaling within the muscle capillary network may mediate this sort of response to increasing metabolism (11), how and where this is sensed, integrated and transmitted through the microvascular network remains to be fully understood. Intriguingly, while the presence of pericytes around capillaries in skeletal muscle has been documented for over 40 years (12), to our knowledge only two recent studies have reported that skeletal muscle pericytes can actively modulate capillary diameter. The first utilised intravital microscopy of the cremaster muscle microcirculation to demonstrate pericyte responsiveness to



endothelin-1 (13). The second study utilised two-photon microscopy to demonstrate that hyperemia-induced changes in capillary diameter in muscle occurred preferentially in pericyte-covered capillary segments and that this vasodilator response was reduced in states of pericyte depletion (14). Based on these emerging evidence from muscle and the ability of pericytes to regulate capillary perfusion in response to increased metabolism in other organs (e.g. the brain) (15), pericytes are ideally positioned between capillaries and skeletal myofibres to be able to sense the metabolic needs of muscle cells and appropriately regulate capillary blood flow.

## **2. Pericytes and capillary blood flow**

The term pericyte was first coined in 1923 (16) and is now used to describe a type of mural cell within the capillary basement membrane. Pericytes encircle capillaries through the extension of cytoplasmic processes that span numerous endothelial cells and often multiple adjacent capillaries (see figure 1); a morphological feature with poorly understood physiological function (17). Of note, pericytes share a number of common features with vascular smooth muscle cells (VSMC) (18) including the cellular machinery necessary for constriction and dilation of blood vessels (19, 20). Pericytes are reported to have a number of functions and can behave as multipotent progenitor cells to facilitate tissue repair including in muscle (21-23). Pericytes have also been reported to actively support angiogenesis (24-26) and capillary function (27) through the release of vascular endothelial growth factor (28). Importantly, brain and retinal pericytes have been shown to contract or relax in response to a number of circulating factors, including angiotensin II and insulin (29, 30), as well as relax in response to cell-metabolism derived factors such as adenosine, lactate and carbon dioxide (CO<sub>2</sub>) (31-33). In the brain, pericytes form an interconnected network within the microvasculature and are capable of transmitting signals upstream to activate adjacent pericytes (34). A recent study calculated that the speed of these conducted responses ranges from 5-20µm/s and that the signal was transmitted both up and downstream of the initiating pericyte (35). This rapid cell-cell communication is particularly intriguing and suggests that rather than acting independently on single capillaries, pericytes may regulate and co-ordinate networks of capillaries through an

integrated and interconnected response to meet tissue metabolic demand. Indeed, we have unpublished histological evidence that this may be the case within the skeletal muscle capillary network (see figure 1). Given their similarity to VSMC's (18), it is also plausible that pericytes may be able to transmit this same signal upstream to dilate or constrict precapillary arterioles to appropriately regulate capillary perfusion. Based on their position at the interface between the capillaries and the cells those capillaries are supplying, pericytes are ideal candidates for the integration of both circulating and local factors to effectively couple blood flow to local metabolism.

## **2.1 Pericytes and capillary blood flow regulation in the brain**

Most of what we know about pericytes stems from studies investigating their role in the brain microcirculation. Indeed, it has been proposed that among all microvascular beds, the brain contains the greatest density of capillary-bound pericytes (36). The intimate association of pericytes with the basement membrane, and complex series of tight junctions and channels linking pericytes and adjacent cells, including endothelial cells, makes their role integral to the structure and function of the blood-brain barrier (37-42). From the abluminal, tissue side, neurons and astrocytes closely associate with brain capillaries at sites immediately adjacent to pericytes (41, 42). This specific co-localisation of cells within the brain microvasculature, termed the neurovascular unit, is the location where nutrient needs of neurons are sensed and integrated into the microvasculature to effect changes in local capillary perfusion; a process called neurovascular coupling. Here, neurotransmitters (e.g. glutamate) and ions (e.g.  $K^+$ ) can be released into the perivascular space where they are sensed by pericytes and endothelial cells to regulate capillary tone and perfusion (37). During periods of increased neural activity, capillary diameter where pericytes reside increases prior to any changes at upstream pre-capillary arterioles (43-45). In response to local glutamate release, pericytes relax and increase capillary diameter through a nitric oxide-dependent activation of the prostaglandin  $E_2$  receptor (15). These were some of the first studies to demonstrate that pericytes are capable of directly affecting capillary diameter and hence blood flow in the brain. Further evidence has recently come from studies investigating the roles that pericytes play in

regulating capillary perfusion in the context of ischemic stroke (15) and Alzheimer's disease (46). Within these pathological settings, pericytes appear to be highly sensitive to ischemia and amyloid  $\beta$  and both cause pericyte constriction that limits capillary blood flow independently of changes in arteriolar tone. Taken together, the above data provide strong evidence for a functional role of pericytes in the tight regulation of capillary blood flow in the brain and their dysregulation in disease states.

## **2.2 Pericytes and capillary blood flow regulation in the heart**

The coronary microcirculation is critical for the normal functioning of the heart, and as with the brain, is acutely affected during periods of ischemia. Despite being the second most abundant cell type in the heart after endothelial cells (47), pericytes retain similar roles as in the brain, including regulation of angiogenesis, barrier potential and selectivity and regulation of capillary blood flow (47). As in the brain, constriction of pericytes has been implicated in the context of myocardial infarction where ischemia causes a powerful constriction to occlude capillaries, which intriguingly can be overcome using intravenous administration of adenosine (47-49). Interestingly, cardiac pericytes may also contribute to the origin of a myocardial infarction through their potential involvement in development of atherosclerosis and vessel stiffening (49). Via a series of pericyte-endothelial cell and pericyte-pericyte gap junctions, cardiac pericytes have been described as forming a functional syncytium capable of propagating electrical and metabolic signals along the vascular tree, akin to the cardiomyocytes themselves (47). Pericytes have also been located within the intima of arterioles and venules of the coronary circulation, appearing to be far more widely distributed along the angioarchitecture with respect to those found in other organs (47). These observations within the heart further implicate the potential role of pericytes as integrators and propagators of metabolic signals where they may not only act locally and directly on capillaries but also signal further upstream to arterioles to modulate total blood flow.

### **2.3 Pericytes and capillary blood flow regulation in the retina**

Due to its developmental origins, the retina is considered a part of the central nervous system (50) and as such the vasculature has similar characteristics to that in the brain. The idea that capillary perfusion is intrinsically linked to photoreceptor activity was first discussed in the 19<sup>th</sup> century (51). Since then, a substantial body of evidence has emerged to suggest that overall increases in total retinal flow (52-54), as well as increases in microvascular flow occur in regions local to photo stimuli (55, 56). Based on pericyte function in the brain (15), investigators have inferred that pericytes may actively regulate retinal capillary blood flow. In contrast, recent experiments performed in the retina by Kornfield and Newman suggest that arterioles, not pericytes are responsible for retinal blood flow regulation in health (57). Despite this, retinal pericyte changes in conjunction with disease have been well documented, particularly in the context of diabetic retinopathy. Retinopathy is a major diabetic complication that leads to vision impairment in 35% of diabetes sufferers world-wide (58) and is characterized by the formation of microaneurysms (59, 60), increased vascular permeability (61) and angiogenesis (58). Together, these microvascular pathologies contribute to hypoxia (62) and vision impairment (63). As with acute hypoxia during ischemic stroke in the brain, chronic hypoxia in the diabetic retina is associated with pericyte dysfunction and a reduced dilatory response that can be reversed by inhibiting inducible nitric oxide synthase (62). While sharing some similar functions to pericytes in the brain and heart, whether retinal pericytes play an active role in regulating retinal capillary flow remains to be definitively determined.

### **2.4 Pericytes and capillary blood flow regulation in the liver**

Despite comprising only ~2-3% of total adult body weight, the liver receives about 25% of total cardiac output at rest (1). To accommodate this high rate of flow, the liver has a unique circulatory system consisting of afferent and efferent vessels connected by specialized exchange vessels termed sinusoids – fenestrated capillaries lacking a basement membrane (64). Compared with capillaries in other organs, fenestrated capillaries facilitate a higher exchange rate between the blood,

endothelial cells and other cells associated with the hepatic sinusoid, including pericytes. Despite these structural capillary differences, hepatic pericytes, alternatively termed Ito cells or hepatic stellate cells, have similar function and morphology to pericytes of other tissues (65, 66). Additionally, they also have a range of liver specific functions including lipid and vitamin A storage (67, 68). Importantly, it has been shown that liver pericytes actively regulate sinusoidal tone in response to locally generated molecules such as endothelin-1 and various prostaglandins (69-73). This normal pericyte response is disrupted in fatty liver disease (70), a condition that involves the accumulation of fat within the liver hepatocytes, causing them to swell and decrease the luminal diameter of surrounding sinusoids (74, 75). Hepatocyte lipid accumulation has been shown to reduce the total number of functional sinusoids (74, 75) and increase pericyte contractility, thus greatly reducing hepatic capillary perfusion (70). Given that microvascular perfusion supports and enables normal liver function, this increased pericyte contractility and hence reduced capillary perfusion is likely to further compromise liver function.

## **2.5 Pericytes and capillary blood flow regulation in the pancreas**

Primarily an exocrine organ, the pancreas contains a small amount of endocrine tissue, only contributing 1-3% of pancreatic volume. Despite this, the endocrine portion has been well studied as it encompasses the islets of Langerhans, housing the pancreatic beta cells responsible for insulin formation and secretion. Pancreatic beta cells are intimately associated with capillaries (76) allowing for prompt assessment of blood glucose concentration and the tailored release of insulin into the blood stream to tightly regulate blood glucose concentrations. Pericytes have been found on capillaries in pancreatic islets (77-79) with evidence suggesting they are necessary for normal beta cell function (78) and regeneration (80). For example, pericytes produce components of the extracellular matrix that promote beta cell proliferation (80), a process that is hindered when pericytes are depleted. Furthermore, it is proposed that pericytes produce growth factors required for glucose-stimulated insulin exocytosis (81). In the absence of these growth factors, beta cells dedifferentiate into an immature form and lose the ability to tailor insulin secretion to blood glucose

concentration (82). Although it appears that pericytes are required for normal insulin secretion, it is still unknown whether pericytes directly communicate with beta cells or whether the majority of these changes are mediated by pericytes manipulating capillary blood flow to beta cells. A recent publication by Almaca et al. demonstrated that pericytes can regulate capillary diameter and blood flow to islets in response to factors such as endothelin-1 (83). The same study went on to show that an increase in beta cell activity resulted in a decrease in pericyte activity, manifesting as capillary dilation (83). Furthermore, increasing concentrations of glucose decreased cytosolic calcium levels in pericytes and simultaneously dilated capillaries (83). This study also highlighted that pericyte coverage of islet capillaries decreases with type 2 diabetes (83). Given that pericytes are necessary for both normal beta cell function and normal islet capillary perfusion, it is plausible that this loss of pericytes on islet capillaries in type 2 diabetes impairs the ability of islet capillary blood flow to be coupled with the metabolic needs of beta cells. Overall, it could be hypothesized that this may compromise both glucose delivery to the beta cells and the subsequent insulin release into the blood stream, thus contributing to progression of type 2 diabetes.

### **3. Can pericytes regulate capillary blood flow in skeletal muscle?**

Due to their substantial contribution to body mass (~40%) and need for energy to drive our movement, mobility and even breathing, skeletal muscles are important determinants of whole-body metabolism. Skeletal muscle resistance vessels supply myofibers via an intricate branching network of arterioles that give rise to terminal pre-capillary arterioles (see figure 1). Each terminal arteriole supplies a group of 15–20 capillaries, collectively known as a microvascular or capillary unit (84). Due to the low metabolic needs of muscle at rest, it is important to note not all parts of muscle capillary networks may be fully perfused at rest (85, 86). Indeed, a number of studies have shown that microvascular perfusion in resting skeletal muscle is neither continuous nor uniform, but rather intermittent and heterogeneous in distribution (87-89). Thus, it has been suggested that microvessels in muscle show rhythmic variations in diameter (vasomotion) and flowmotion to alternate blood flow through different capillary units as needed to maintain resting metabolism (86).

When metabolism increases, for example during muscle contraction, dilation of the vasculature enhances the delivery of oxygen and nutrients to active muscle fibres in concert with removing metabolic waste products (90), thereby coupling local perfusion to metabolic demand. The extent to which capillary blood flow increases during muscle contraction is related intimately to the power and force of contraction (and hence metabolic demand) (91). Importantly, blood flow is increased only to active muscles and can be specifically directed only to active fibres within a muscle (92, 93). This is reminiscent of the blood flow response in the brain (as discussed above) and suggests that the muscle microvasculature may be able to respond to highly localised metabolic changes and demands.

Given that skeletal muscle capillary units are supplied by terminal arterioles, the majority of research thus far has focused on arteriolar responses to increased metabolic demand. Indeed, numerous of studies have shown that arterioles promptly dilate in response to exercise (85, 94, 95), metabolites such as adenosine (96, 97) and prostaglandins (98, 99). Given the evidence summarized above and the anatomical organization of the muscle vasculature, it stands to reason that arteriolar constriction/dilation is a major determinant of vascular resistance and hence capillary perfusion in muscle during exercise (5, 10). However, given the presence of pericytes around skeletal muscle capillaries, and the emerging evidence for pericyte contraction and dilation (13, 14), capillaries (via pericyte actions) are able to alter capillary blood flow and/or blood content independently from, or in conjunction with arterioles to determine nutrient delivery to skeletal muscle fibres to control local metabolism (100). While these subtle capillary flow oscillations may not be easily distinguishable during high intensity exercise, they may be identifiable at lower work rates that do not increase bulk flow or in instances where muscle capillary perfusion is increased independent of muscle contraction (100). Whether pericytes participate in or regulate capillary perfusion in these contexts remains to be determined.

As well as facilitating contraction and movement, skeletal muscles are also a major site for insulin-mediated glucose disposal and contribute greatly to whole body metabolism. For this reason, blunted insulin-mediated muscle glucose uptake has been championed as one of the early and

primary drivers of development and progression of whole body insulin resistance (101). Within muscle, an important action of insulin is to first increase capillary blood flow (102). In doing so, insulin enhances the delivery of both glucose and insulin to muscle cells to facilitate insulin-stimulated muscle glucose uptake. While local infusion of insulin into the skeletal muscle microvasculature stimulates increased capillary blood flow and enhances glucose uptake (103), central nervous system administration of insulin does not increase capillary perfusion nor glucose uptake in muscle (104). These data would indicate that insulin's action to increase capillary blood flow within muscle is a localised response and mediated through direct effects on the microvasculature. In addition to this, the microvasculature is one of the first tissues to lose its response to insulin during the development of insulin resistance and this is closely associated with reduced insulin-mediated glucose disposal (105-108). Thus, while it has been well established that insulin acts to increase capillary perfusion, identifying exactly where and how insulin exerts its effects in the vascular tree has remained elusive. Importantly, studies have shown that insulin mediates its vascular actions through nitric oxide and endothelin-1 (109-118). Moreover, Spiranac et al. recently reported that endothelin-1 provokes strong and stable contraction of individual capillary pericytes in mouse cremaster muscle in vivo via a  $\text{Ca}^{2+}$ -mediated mechanism (13). Coupled with the knowledge that muscle capillaries are almost completely covered by pericytes (figure 1C), that insulin can act directly on pericytes in other organs (30), and that pericytes are lost from skeletal muscle capillaries in type 2 diabetes (119), it is tempting to speculate that pericytes may contribute to insulin's actions on the muscle microvasculature and that these may be lost in insulin resistance. Whether this hypothesis is correct is currently not known and warrants future investigation.

### **Conclusions and future directions**

In this review, we have highlighted the emerging evidence for pericyte-mediated regulation of capillary blood flow across various organs within the body. While the concept that pericytes actively regulate capillary blood flow in muscle is in its relative infancy, current studies suggest that (i) pericytes are present around skeletal muscle capillaries (figure 1; (13, 14)); (ii) skeletal muscle pericytes have the capacity to regulate capillary diameter (13, 14) and (iii) degeneration of capillary



pericytes occurs in skeletal muscle during type 2 diabetes (119). Based on the emerging repertoire of pericyte functions in other organs, the immediate need in skeletal muscle research is to (i) determine whether skeletal muscle pericytes are able to actively regulate capillary tone in response to circulating and metabolic factors; (ii) determine how pericyte loss from the microvasculature impacts muscle function and microvascular responses; and (iii) characterise the anatomical locations of pericytes in skeletal muscle and determine whether this differs between different muscle fibre types. Based on what is known about pericyte function in other organs and the emerging evidence in muscle, we hypothesize that pericytes act as gatekeepers, controlling muscle nutrient exchange, coupling metabolism to microvascular supply (see figure 2 for working hypothesis). Given the strategic location of pericytes between capillaries and muscle fibres, unravelling their functions in skeletal muscle will significantly advance our understanding of muscle physiology during health, exercise and in disease states such as insulin resistance and exercise intolerance.

### **Acknowledgements**

This work was supported by funding from the University of Tasmania, the Royal Hobart Hospital Research Foundation and the National Health and Medical Research Council (NHMRC; APP1137776). The authors declare no conflict of interest.

### **REFERENCES**

1. Williams L, Leggett R. Reference values for resting blood flow to organs of man. *Clinical Physics and Physiological Measurement*. 1989;10(3):187.
2. Joyner MJ, Casey DP. Regulation of increased blood flow (hyperemia) to muscles during exercise: a hierarchy of competing physiological needs. *Physiological reviews*. 2015;95(2):549-601.
3. Krogh A. The supply of oxygen to the tissues and the regulation of the capillary circulation. *The Journal of Physiology*. 1919;52(6):457-74.

4. Premilovac D, Attrill E, Rattigan S, Richards SM, Kim J-, Keske MA. Acute, local infusion of angiotensin II impairs microvascular and metabolic insulin sensitivity in skeletal muscle. *Cardiovascular research*. 2018.
5. Segal SS. Regulation of blood flow in the microcirculation. *Microcirculation*. 2005;12(1):33-45.
6. Bagher P, Segal SS. Regulation of blood flow in the microcirculation: role of conducted vasodilation. *Acta Physiologica*. 2011;202(3):271-84.
7. Sinkler SY, Segal SS. Rapid versus slow ascending vasodilatation: intercellular conduction versus flow - mediated signalling with tetanic versus rhythmic muscle contractions. *The Journal of physiology*. 2017;595(23):7149-65.
8. Yu J, Bihari A, Lidington D, Tyml K. Gap junction uncouplers attenuate arteriolar response to distal capillary stimuli. *Microvascular research*. 2000;59(1):162-8.
9. Berg BR, Cohen KD, Sarelius IH. Direct coupling between blood flow and metabolism at the capillary level in striated muscle. *American Journal of Physiology-Heart and Circulatory Physiology*. 1997;272(6):H2693-H700.
10. Cohen KD, Berg BR, Sarelius IH. Remote arteriolar dilations in response to muscle contraction under capillaries. *American Journal of Physiology-Heart and Circulatory Physiology*. 2000;278(6):H1916-H23.
11. Novielli NM, Murrant CL. Propagation of Capillary-initiated Conducted Vasodilation in Skeletal Muscle—A Novel Paracrine Signaling Pathway. *The FASEB Journal*. 2016;30(1\_supplement):945.17-.17.
12. Tilton RG, Kilo C, Williamson JR, Murch DW. Differences in pericyte contractile function in rat cardiac and skeletal muscle microvasculatures. *Microvascular research*. 1979;18(3):336-52.
13. Špiranec K, Chen W, Werner F, Nikolaev VO, Naruke T, Koch F, et al. Endothelial C-type natriuretic peptide acts on pericytes to regulate microcirculatory flow and blood pressure. *Circulation*. 2018;138(5):494-508.
14. Methner C, Mishra A, Golgotiu K, Li Y, Wei W, Yanez ND, et al. Pericyte constriction underlies capillary de-recruitment during hyperemia in the setting of arterial stenosis. *American Journal of Physiology-Heart and Circulatory Physiology*. 2019.
15. Hall CN, Reynell C, Gesslein B, Hamilton NB, Mishra A, Sutherland BA, et al. Capillary pericytes regulate cerebral blood flow in health and disease. *Nature*. 2014;508(7494):55-60.
16. Zimmermann KW. Der feinere bau der blutcapillaren. *Zeitschrift für Anatomie und Entwicklungsgeschichte*. 1923;68(1):29-109.

17. Attwell D, Mishra A, Hall CN, O'Farrell FM, Dalkara T. What is a pericyte? *Journal of Cerebral Blood Flow & Metabolism*. 2016;36(2):451-5.
18. Smyth LC, Rustenhoven J, Scotter EL, Schweder P, Faull RL, Park TI, et al. Markers for human brain pericytes and smooth muscle cells. *Journal of chemical neuroanatomy*. 2018;92:48-60.
19. Yamanishi S, Katsumura K, Kobayashi T, Puro DG. Extracellular lactate as a dynamic vasoactive signal in the rat retinal microvasculature. *American Journal of Physiology-Heart and Circulatory Physiology*. 2006;290(3):H925-H34.
20. Herman IM, D'amore PA. Microvascular pericytes contain muscle and nonmuscle actins. *The Journal of cell biology*. 1985;101(1):43-52.
21. Dellavalle A, Maroli G, Covarello D, Azzoni E, Innocenzi A, Perani L, et al. Pericytes resident in postnatal skeletal muscle differentiate into muscle fibres and generate satellite cells. *Nature communications*. 2011;2:499.
22. Dellavalle A, Sampaolesi M, Tonlorenzi R, Tagliafico E, Sacchetti B, Perani L, et al. Pericytes of human skeletal muscle are myogenic precursors distinct from satellite cells. *Nature cell biology*. 2007;9(3):255.
23. Asahina K, Sato H, Yamasaki C, Kataoka M, Shiokawa M, Katayama S, et al. Pleiotrophin/heparin-binding growth-associated molecule as a mitogen of rat hepatocytes and its role in regeneration and development of liver. *The American journal of pathology*. 2002;160(6):2191-205.
24. Ogura S, Kurata K, Hattori Y, Takase H, Ishiguro-Oonuma T, Hwang Y, et al. Sustained inflammation after pericyte depletion induces irreversible blood-retina barrier breakdown. *JCI insight*. 2017;2(3).
25. Smith AO, Bowers SL, Stratman AN, Davis GE. Hematopoietic stem cell cytokines and fibroblast growth factor-2 stimulate human endothelial cell-pericyte tube co-assembly in 3D fibrin matrices under serum-free defined conditions. *PLoS One*. 2013;8(12):e85147.
26. Stratman AN, Malotte KM, Mahan RD, Davis MJ, Davis GE. Pericyte recruitment during vasculogenic tube assembly stimulates endothelial basement membrane matrix formation. *Blood*. 2009;114(24):5091-101.
27. Bell RD, Winkler EA, Sagare AP, Singh I, LaRue B, Deane R, et al. Pericytes control key neurovascular functions and neuronal phenotype in the adult brain and during brain aging. *Neuron*. 2010;68(3):409-27.
28. Mancuso MR, Davis R, Norberg SM, O'Brien S, Sennino B, Nakahara T, et al. Rapid vascular regrowth in tumors after reversal of VEGF inhibition. *The Journal of clinical investigation*. 2006;116(10):2610-21.
29. Dodge AB, Hechtman HB, Shepro D. Microvascular endothelial-derived autacoids regulate pericyte contractility. *Cell motility and the cytoskeleton*. 1991;18(3):180-8.

30. King GL, Buzney SM, Kahn CR, Hetu N, Buchwald S, Macdonald S, et al. Differential responsiveness to insulin of endothelial and support cells from micro-and macrovessels. *The Journal of clinical investigation*. 1983;71(4):974-9.
31. Chen Q, Anderson DR. Effect of CO<sub>2</sub> on intracellular pH and contraction of retinal capillary pericytes. *Investigative ophthalmology & visual science*. 1997;38(3):643-51.
32. Dirnagl U, Niwa K, Lindauer U, Villringer A. Coupling of cerebral blood flow to neuronal activation: role of adenosine and nitric oxide. *American Journal of Physiology-Heart and Circulatory Physiology*. 1994;267(1):H296-H301.
33. Neuhaus AA, Sutherland BA, Buchan AM. Targeting Pericytes and the Microcirculation for Ischemic Stroke Therapy. *Neuroprotective Therapy for Stroke and Ischemic Disease: Springer*; 2017. p. 537-56.
34. Peppiatt CM, Howarth C, Mobbs P, Attwell D. Bidirectional control of CNS capillary diameter by pericytes. *Nature*. 2006;443(7112):700.
35. Cai C, Fordsmann JC, Jensen SH, Gesslein B, Lønstrup M, Hald BO, et al. Stimulation-induced increases in cerebral blood flow and local capillary vasoconstriction depend on conducted vascular responses. *Proceedings of the National Academy of Sciences*. 2018;115(25):E5796-E804.
36. Santos GS, Magno LA, Romano-Silva MA, Mintz A, Birbrair A. Pericyte Plasticity in the Brain. *Neuroscience bulletin*. 2019;35(3):551-60.
37. Attwell D, Buchan AM, Charkpak S, Lauritzen M, MacVicar BA, Newman EA. Glial and neuronal control of brain blood flow. *Nature*. 2010;468(7321):232.
38. Busija DW, Bari F, Domoki F, Louis T. Mechanisms involved in the cerebrovascular dilator effects of N-methyl-d-aspartate in cerebral cortex. *Brain research reviews*. 2007;56(1):89-100.
39. Longden TA, Hill-Eubanks DC, Nelson MT. Ion channel networks in the control of cerebral blood flow. *Journal of Cerebral Blood Flow & Metabolism*. 2016;36(3):492-512.
40. Muoio V, Persson P, Sendeski M. The neurovascular unit—concept review. *Acta physiologica*. 2014;210(4):790-8.
41. Sweeney MD, Ayyadurai S, Zlokovic BV. Pericytes of the neurovascular unit: key functions and signaling pathways. *Nature neuroscience*. 2016;19(6):771.
42. McConnell HL, Kersch CN, Woltjer RL, Neuwelt EA. The translational significance of the neurovascular unit. *Journal of Biological Chemistry*. 2017;292(3):762-70.

43. Cheng J, Korte N, Nortley R, Sethi H, Tang Y, Attwell D. Targeting pericytes for therapeutic approaches to neurological disorders. *Acta Neuropathol.* 2018;136(4):507-23.
44. Kisler K, Nelson AR, Rege SV, Ramanathan A, Wang Y, Ahuja A, et al. Pericyte degeneration leads to neurovascular uncoupling and limits oxygen supply to brain. *Nat Neurosci.* 2017;20(3):406-16.
45. Mishra A, Reynolds JP, Chen Y, Gourine AV, Rusakov DA, Attwell D. Astrocytes mediate neurovascular signaling to capillary pericytes but not to arterioles. *Nat Neurosci.* 2016;19(12):1619-27.
46. Nortley R, Korte N, Izquierdo P, Hirunpattarasilp C, Mishra A, Jaunmuktane Z, et al. Amyloid  $\beta$  oligomers constrict human capillaries in Alzheimer's disease via signaling to pericytes. *Science.* 2019:eaav9518.
47. Nees S, Weiss DR, Juchem G. Focus on cardiac pericytes. *Pflugers Arch.* 2013;465(6):779-87.
48. Costa MA, Paiva AE, Andreotti JP, Cardoso MV, Cardoso CD, Mintz A, et al. Pericytes constrict blood vessels after myocardial ischemia. *J Mol Cell Cardiol.* 2018;116:1-4.
49. O'Farrell FM, Mastitskaya S, Hammond-Haley M, Freitas F, Wah WR, Attwell D. Capillary pericytes mediate coronary no-reflow after myocardial ischaemia. *Elife.* 2017;6.
50. Etchevers HC, Vincent C, Le Douarin NM, Couly GF. The cephalic neural crest provides pericytes and smooth muscle cells to all blood vessels of the face and forebrain. *Development.* 2001;128(7):1059-68.
51. Roy CS, Sherrington CS. On the regulation of the blood-supply of the brain. *The Journal of physiology.* 1890;11(1-2):85-158.
52. Falsini B, Riva CE, Logean E. Flicker-evoked changes in human optic nerve blood flow: relationship with retinal neural activity. *Investigative ophthalmology & visual science.* 2002;43(7):2309-16.
53. Formaz F, Riva C, Geiser M. Diffuse luminance flicker increases retinal vessel diameter in humans. *Current eye research.* 1997;16(12):1252-7.
54. Garhöfer G, Zawinka C, Resch H, Kothy P, Schmetterer L, Dorner G. Reduced response of retinal vessel diameters to flicker stimulation in patients with diabetes. *British Journal of Ophthalmology.* 2004;88(7):887-91.
55. Duong TQ, Ngan S-C, Ugurbil K, Kim S-G. Functional magnetic resonance imaging of the retina. *Investigative ophthalmology & visual science.* 2002;43(4):1176-81.
56. Srienc AI, Kurth-Nelson ZL, Newman E. Imaging retinal blood flow with laser speckle flowmetry. *Frontiers in neuroenergetics.* 2010;2:128.

57. Kornfield TE, Newman EA. Regulation of blood flow in the retinal trilaminar vascular network. *Journal of Neuroscience*. 2014;34(34):11504-13.
58. Hammes H-P. Diabetic retinopathy: hyperglycaemia, oxidative stress and beyond. *Diabetologia*. 2018;61(1):29-38.
59. Cogan DG, Toussaint D, Kuwabara T. Retinal vascular patterns: IV. Diabetic retinopathy. *Archives of Ophthalmology*. 1961;66(3):366-78.
60. Lindahl P, Johansson BR, Levéen P, Betsholtz C. Pericyte loss and microaneurysm formation in PDGF-B-deficient mice. *Science*. 1997;277(5323):242-5.
61. Ben-nun J, Alder VA, Constable IJ. Retinal microvascular patency in the diabetic rat. *International ophthalmology*. 2004;25(4):187-92.
62. Hammes H-P, Lin J, Renner O, Shani M, Lundqvist A, Betsholtz C, et al. Pericytes and the pathogenesis of diabetic retinopathy. *Diabetes*. 2002;51(10):3107-12.
63. Yau J, Rogers SL, Kawasaki R, Lamoureux EL, Kowalski JW, Bek T, et al. Meta-Analysis for Eye Disease (META-EYE) Study Group. Global prevalence and major risk factors of diabetic retinopathy. *Diabetes care*. 2012;35(3):556-64.
64. Wisse E, De Zanger R, Charels K, Van Der Smissem P, McCuskey R. The liver sieve: considerations concerning the structure and function of endothelial fenestrae, the sinusoidal wall and the space of Disse. *Hepatology*. 1985;5(4):683-92.
65. Ito T NM. Kupfer's cells and fat storing cells in the capillary wall of human liver. *Okajimas Folia Anat Jpn*. 1952;24(4):243-58.
66. Ahern M, Hall P, Halliday J, Liddle C, Olynyk J, Ramm G, et al. Hepatic stellate cell nomenclature. *Hepatology*. 1996;23(1):193-.
67. Blomhoff R, Rasmussen M, Nilsson A, Norum K, Berg T, Blaner W, et al. Hepatic retinol metabolism. Distribution of retinoids, enzymes, and binding proteins in isolated rat liver cells. *Journal of Biological Chemistry*. 1985;260(25):13560-5.
68. Senoo H. Structure and function of hepatic stellate cells. *Medical Electron Microscopy*. 2004;37(1):3-15.
69. Bauer M, Bauer I, Sonin NV, Kresge N, Baveja R, Yokoyama Y, et al. Functional significance of endothelin B receptors in mediating sinusoidal and extrasinusoidal effects of endothelins in the intact rat liver. *Hepatology*. 2000;31(4):937-47.

70. Bauer M, Paquette NC, Zhang JX, Bauer I, Pannen BH, Kleeberger SR, et al. Chronic ethanol consumption increases hepatic sinusoidal contractile response to endothelin - 1 in the rat. *Hepatology*. 1995;22(5):1565-76.
71. Oda M, Han JY, Yokomori H. Local regulators of hepatic sinusoidal microcirculation: recent advances. *Clinical hemorheology and microcirculation*. 2000;23(2, 3, 4):85-94.
72. Pinzani M. Novel insights into the biology and physiology of the Ito cell. *Pharmacology & therapeutics*. 1995;66(2):387-412.
73. Thimigan MS, Yee Jr HF. Quantitation of rat hepatic stellate cell contraction: stellate cells' contribution to sinusoidal resistance. *American Journal of Physiology-Gastrointestinal and Liver Physiology*. 1999;277(1):G137-G43.
74. Sun C-K, Zhang X-Y, Wheatley AM. Increased NAD (P) H fluorescence with decreased blood flow in the steatotic liver of the obese Zucker rat. *Microvascular research*. 2003;66(1):15-21.
75. Sun C-K, Zhang X-Y, Zimmermann A, Davis G, Wheatley AM. Effect of ischemia-reperfusion injury on the microcirculation of the steatotic liver of the Zucker Rat1. *Transplantation*. 2001;72(10):1625-31.
76. Richards OC, Raines SM, Attie AD. The role of blood vessels, endothelial cells, and vascular pericytes in insulin secretion and peripheral insulin action. *Endocrine reviews*. 2010;31(3):343-63.
77. Armulik A, Genové G, Betsholtz C. Pericytes: developmental, physiological, and pathological perspectives, problems, and promises. *Developmental cell*. 2011;21(2):193-215.
78. Sasson A, Rach E, Sakhneny L, Baer D, Lisnyansky M, Epshtein A, et al. Islet pericytes are required for  $\beta$ -cell maturity. *Diabetes*. 2016;65(10):3008-14.
79. Tang S-C, Chiu Y-C, Hsu C-T, Peng S-J, Fu Y-Y. Plasticity of Schwann cells and pericytes in response to islet injury in mice. *Diabetologia*. 2013;56(11):2424-34.
80. Epshtein A, Rach E, Sakhneny L, Mizrachi S, Baer D, Landsman L. Neonatal pancreatic pericytes support  $\beta$ -cell proliferation. *Molecular metabolism*. 2017;6(10):1330-8.
81. Houtz J, Borden P, Ceasrine A, Minichiello L, Kuruvilla R. Neurotrophin signaling is required for glucose-induced insulin secretion. *Developmental cell*. 2016;39(3):329-45.
82. Talchai C, Xuan S, Lin HV, Sussel L, Accili D. Pancreatic  $\beta$  cell dedifferentiation as a mechanism of diabetic  $\beta$  cell failure. *Cell*. 2012;150(6):1223-34.
83. Almaca J, Weitz J, Rodriguez-Diaz R, Pereira E, Caicedo A. The Pericyte of the Pancreatic Islet Regulates Capillary Diameter and Local Blood Flow. *Cell Metabolism*. 2018;27(3):630-44.e4.

84. Delashaw JB, Duling BR. A study of the functional elements regulating capillary perfusion in striated muscle. *Microvascular Research*. 1988;36:162-71.
85. Honig CR, Odoroff CL, Frierson JL. Active and passive capillary control in red muscle at rest and in exercise. *American Journal of Physiology*. 1982;243:H196-H206.
86. Murrant CL, Sarelius IH. Local control of blood flow during active hyperaemia: what kinds of integration are important? *The Journal of physiology*. 2015;593(21):4699-711.
87. Lindbom L, Arfors KE. Non-homogeneous blood flow distribution in the rabbit tenuissimus muscle. Differential control of total blood flow and capillary perfusion. *Acta Physiologica Scandinavica*. 1984;122:225-33.
88. Piiper J, Marconi C, Heisler N, Meyer M, Weitz H, Pendergast DR, et al. Spatial and temporal variability of blood flow in stimulated dog gastrocnemius muscle. *Advances in Experimental Medicine and Biology*. 1989;248:719-43.
89. Blackwood SJ, Dwyer RM, Bradley EA, Keske MA, Richards SM, Rattigan S. Determination of skeletal muscle microvascular flowmotion with contrast-enhanced ultrasound. *Ultrasound in Medicine and Biology*. 2017;43(9):2013-23.
90. VanTeeffelen JW, Segal SS. Effect of motor unit recruitment on functional vasodilatation in hamster retractor muscle. *Journal of Physiology*. 2000;524(1):267-78.
91. Radegran G, Saltin B. Muscle blood flow at onset of dynamic exercise in humans. *American Journal of Physiology-Heart and Circulatory Physiology*. 1998;274(1):H314-H22.
92. Mackie BG, Terjung RL. Blood flow to different skeletal muscle fiber types during contraction. *American Journal of Physiology-Heart and Circulatory Physiology*. 1983;245(2):H265-H75.
93. Twynstra J, Ruiz D, Murrant C. Functional coordination of the spread of vasodilations through skeletal muscle microvasculature: implications for blood flow control. *Acta Physiologica*. 2012;206(4):229-41.
94. VanTeeffelen JW, Segal SS. Rapid dilation of arterioles with single contraction of hamster skeletal muscle. *American Journal of Physiology-Heart and Circulatory Physiology*. 2006;290(1):H119-H27.
95. Honig CR, Odoroff CL, Frierson JL. Capillary recruitment in exercise: rate, extent, uniformity, and relation to blood flow. *American Journal of Physiology-Heart and Circulatory Physiology*. 1980;238(1):H31-H42.



96. Murrant CL, Sarelius IH. Multiple dilator pathways in skeletal muscle contraction-induced arteriolar dilations. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*. 2002;282(4):R969-R78.
97. Poucher S, Nowell C, Collis M. The role of adenosine in exercise hyperaemia of the gracilis muscle in anaesthetized cats. *The journal of physiology*. 1990;427(1):19-29.
98. Koller A, Bagi Z. On the role of mechanosensitive mechanisms eliciting reactive hyperemia. *American Journal of Physiology-Heart and Circulatory Physiology*. 2002;283(6):H2250-H9.
99. Koeppen M, Feil R, Siegl D, Feil S, Hofmann F, Pohl U, et al. cGMP-dependent protein kinase mediates NO-but not acetylcholine-induced dilations in resistance vessels in vivo. *Hypertension*. 2004;44(6):952-5.
100. Inyard AC, Clerk LH, Vincent MA, Barrett EJ. Contraction stimulates nitric oxide-independent microvascular recruitment and increases muscle insulin uptake. *Diabetes*. 2007;56(9):2194-200.
101. DeFronzo RA, Tripathy D. Skeletal muscle insulin resistance is the primary defect in type 2 diabetes. *Diabetes care*. 2009;32(suppl 2):S157-S63.
102. Keske MA, Premilovac D, Bradley EA, Dwyer RM, Richards SM, Rattigan S. Muscle microvascular blood flow responses in insulin resistance and ageing. *The Journal of physiology*. 2016;594(8):2223-31.
103. Coggins M, Lindner J, Rattigan S, Jahn L, Fasy E, Kaul S, et al. Physiologic hyperinsulinemia enhances human skeletal muscle perfusion by capillary recruitment. *Diabetes*. 2001;50(12):2682-90.
104. Bradley E, Willson K, Choi-Lundberg D, Clark M, Rattigan S. Effects of central administration of insulin or I - NMMA on rat skeletal muscle microvascular perfusion. *Diabetes, Obesity and Metabolism*. 2010;12(10):900-8.
105. Bonner JS, Lantier L, Hasenour CM, James FD, Bracy DP, Wasserman DH. Muscle-specific vascular endothelial growth factor deletion induces muscle capillary rarefaction creating muscle insulin resistance. *Diabetes*. 2013;62(2):572-80.
106. Kubota T, Kubota N, Kumagai H, Yamaguchi S, Kozono H, Takahashi T, et al. Impaired insulin signaling in endothelial cells reduces insulin-induced glucose uptake by skeletal muscle. *Cell metabolism*. 2011;13(3):294-307.
107. Premilovac D, Bradley EA, Ng HL, Richards SM, Rattigan S, Keske MA. Muscle insulin resistance resulting from impaired microvascular insulin sensitivity in Sprague Dawley rats. *Cardiovascular research*. 2013;98(1):28-36.


108. Premilovac D, Richards SM, Rattigan S, Keske MA. A vascular mechanism for high-sodium-induced insulin resistance in rats. *Diabetologia*. 2014;57(12):2586-95.
109. Montagnani M, Ravichandran LV, Chen H, Esposito DL, Quon MJ. Insulin receptor substrate-1 and phosphoinositide-dependent kinase-1 are required for insulin-stimulated production of nitric oxide in endothelial cells. *Molecular endocrinology*. 2002;16(8):1931-42.
110. Steinberg HO, Brechtel G, Johnson A, Fineberg N, Baron AD. Insulin-mediated skeletal muscle vasodilation is nitric oxide dependent. A novel action of insulin to increase nitric oxide release. *The Journal of clinical investigation*. 1994;94(3):1172-9.
111. Vincent M, Barrett E, Lindner J, Clark M, Rattigan S. Inhibiting NOS blocks microvascular recruitment and blunts muscle glucose uptake in response to insulin. *American Journal of Physiology-Endocrinology And Metabolism*. 2003;285(1):E123-E9.
112. Cardillo C, Campia U, Bryant MB, Panza JA. Increased activity of endogenous endothelin in patients with type II diabetes mellitus. *Circulation*. 2002;106(14):1783-7.
113. Cardillo C, Nambi SS, Kilcoyne CM, Choucair WK, Katz A, Quon MJ, et al. Insulin stimulates both endothelin and nitric oxide activity in the human forearm. *Circulation*. 1999;100(8):820-5.
114. Potenza MA, Marasciulo FL, Chieppa DM, Brigiani GS, Formoso G, Quon MJ, et al. Insulin resistance in spontaneously hypertensive rats is associated with endothelial dysfunction characterized by imbalance between NO and ET-1 production. *American Journal of Physiology-Heart and Circulatory Physiology*. 2005;289(2):H813-H22.
115. Ross R, Kolka C, Rattigan S, Clark M. Acute blockade by endothelin-1 of haemodynamic insulin action in rats. *Diabetologia*. 2007;50(2):443-51.
116. Zeng G, Nystrom FH, Ravichandran LV, Cong L-N, Kirby M, Mostowski H, et al. Roles for insulin receptor, PI3-kinase, and Akt in insulin-signaling pathways related to production of nitric oxide in human vascular endothelial cells. *Circulation*. 2000;101(13):1539-45.
117. Zeng G, Quon MJ. Insulin-stimulated production of nitric oxide is inhibited by wortmannin. Direct measurement in vascular endothelial cells. *The Journal of clinical investigation*. 1996;98(4):894-8.
118. Eringa EC, Stehouwer CD, Walburg K, Clark AD, van Nieuw Amerongen GP, Westerhof N, et al. Physiological concentrations of insulin induce endothelin-dependent vasoconstriction of skeletal muscle resistance arteries in the presence of tumor necrosis factor- $\alpha$  dependence on c-Jun N-terminal kinase. *Arteriosclerosis, thrombosis, and vascular biology*. 2006;26(2):274-80.

119. Tilton RG, Hoffmann PL, Kilo C, Williamson JR. Pericyte degeneration and basement membrane thickening in skeletal muscle capillaries of human diabetics. *Diabetes*. 1981;30(4):326-34.


### Figure legends

**Figure 1. Skeletal muscle capillaries are covered by pericytes.** Diagram depicting the structure of the microvasculature in skeletal muscle (**A**) highlighting the transition from pre-capillary arterioles covered by vascular smooth muscle cells (VSMCs) to capillaries covered almost completely by pericytes (**B** and **C**). In panel **C**, we performed immunohistochemistry on longitudinal sections of rat tibialis anterior muscle to assess pericyte (red; anti-NG2 antibody; Merck Millipore Cat.No. AB5320) coverage of capillaries (green; tomato lectin; Vector Labs; Cat.No. DL-1174). Nuclei were stained with DAPI (4',6-diamidino-2-phenylindole) and are shown in blue. White arrows point to pericyte cell bodies; the yellow arrow highlights a pericyte process extending across a muscle fibre to wrap around two adjacent capillaries. Scale bar = 25µm.

**Figure 2. Working hypothesis where pericytes dilate capillaries to regulate blood flow and skeletal muscle metabolism.** **1.** Circulating factors such as insulin (I; shown in green) act on endothelial cells to stimulate nitric oxide (NO) production by endothelial nitric oxide synthase (eNOS). NO diffuses from endothelial cells and stimulates pericyte relaxation. **2.** Insulin transitions from the capillary lumen into the interstitial space through either transcellular or paracellular pathways and directly acts on pericytes to stimulate potassium (K<sup>+</sup>)-dependent hyperpolarization and relaxation. **3.** Muscle metabolism derived molecules, including carbon dioxide (CO<sub>2</sub>), lactate (Lac) and adenosine (Ad), diffuse from muscle cells and act on pericytes to cause relaxation. **4.** In each case above, pericyte relaxation causes an increase in capillary diameter, enhancing delivery of glucose for storage as glycogen in the presence of insulin or to facilitate muscle metabolism.



cep\_13208\_f1.jpg



cep\_13208\_f2.jpg